

Measurement of the absolute branching fraction of the $D_s^\pm$ meson.

K. Abe, I. Adachi, J. Dragic, H. Fujii, T. Gershon, J. Haba, M. Hazumi, T. Higuchi,
 Y. Igarashi, R. Itoh, Y. Iwasaki, N. Katayama, H. Kichimi, P. Krokovny,
 A. Limosani, I. Nakamura, M. Nakao, H. Nakazawa, S. Nishida, T. Nozaki,
 H. Ozaki, F. J. Ronga, S. Saitoh, Y. Sakai, R. Stamen, K. Sumisawa, S. Y. Suzuki,
 O. Tajima, F. Takasaki, K. Tamai, M. Tanaka, K. Trabelsi, T. Tsuboyama,
 T. Tsukamoto, S. Uehara, Y. Unno, S. Uno, Y. Ushiroda, M. Yamauchi, and J. Zhang
High Energy Accelerator Research Organization (KEK), Tsukuba

K. Abe, Y. Hoshi, and K. Neichi
Tohoku Gakuin University, Tagajo

H. Aihara, N. C. Hastings, A. Ishikawa, K. Itoh, M. Iwasaki,
 H. Kakuno, A. Kusaka, Y. Nakahama, and K. Tanabe
Department of Physics, University of Tokyo, Tokyo

D. Anipko, K. Arinstein, V. Aulchenko, I. Bedny, A. Bondar,
 S. Eidelman, D. Epifanov, N. Gabyshev, A. Kuzmin, A. Poluektov,
 N. Root, B. Shwartz, V. Sidorov, Y. Usov, and V. Zhilich
Budker Institute of Nuclear Physics, Novosibirsk

K. Aoki, Y. Enari, K. Hara, K. Hayasaka, T. Hokuue, T. Iijima,
 K. Ikado, K. Inami, N. Kishimoto, Y. Kozakai, T. Kubota, Y. Miyazaki,
 T. Ohshima, T. Okabe, N. Sato, K. Senyo, and S. Yoshino
Nagoya University, Nagoya

T. Arakawa, T. Kawasaki, H. Miyata, N. Tamura, and M. Watanabe
Niigata University, Niigata

Y. Asano
University of Tsukuba, Tsukuba

T. Aso

Toyama National College of Maritime Technology, Toyama

T. Aushev, A. Bay, L. Hinz, C. Jacoby, T. Schietinger,

O. Schneider, S. Villa, J. Wicht, and D. Zürcher

Swiss Federal Institute of Technology of Lausanne, EPFL, Lausanne

T. Aziz, S. Banerjee, G. Gokhroo, and G. Majumder

Tata Institute of Fundamental Research, Bombay

S. Bahinipati, A. Drutskoy, P. Goldenzweig, K. Kinoshita,

R. Kulasiri, K. Sayeed, A. J. Schwartz, and A. Somov

University of Cincinnati, Cincinnati, Ohio 45221

A. M. Bakich, S. Cole, S. McOnie, N. Parslow,

L. S. Peak, H. Stoeck, K. E. Varvell, and B. D. Yabsley

University of Sydney, Sydney NSW

V. Balagura, R. Chistov, M. Danilov, D. Liventsev, T. Medvedeva,

R. Mizuk, P. Pakhlov, G. Pakhlova, I. Tikhomirov, and T. Uglov

Institute for Theoretical and Experimental Physics, Moscow

Y. Ban and X. C. Tian

Peking University, Beijing

E. Barberio, J. Dalseno, R. Dowd, G. R. Moloney,

M. E. Sevier, G. N. Taylor, Y. F. Tse, and P. Urquijo

University of Melbourne, Victoria

M. Barbero, T. E. Browder, H. Guler, M. Jones, J. Li, K. Nishimura,

S. L. Olsen, M. Peters, J. Rorie, H. Sahoo, K. Uchida, and G. Varner

University of Hawaii, Honolulu, Hawaii 96822

K. Belous, M. Shapkin, and A. Sokolov

Institute of High Energy Physics, Protvino

U. Bitenc, I. Bizjak, S. Fratina, A. Gorišek, R. Pestotnik, M. Starič, and A. Zupanc

J. Stefan Institute, Ljubljana

S. Blyth, A. Chen, W. T. Chen, A. Go, S. Hou, and C. C. Kuo

National Central University, Chung-li

A. Bozek, P. Kapusta, T. Lesiak, A. Matyja, Z. Natkaniec,
W. Ostrowicz, H. Palka, M. Rozanska, and J. Wiechczynski

H. Niewodniczanski Institute of Nuclear Physics, Krakow

M. Bračko and S. Korpar

University of Maribor, Maribor and

J. Stefan Institute, Ljubljana

J. Brodzicka

High Energy Accelerator Research Organization (KEK), Tsukuba and

H. Niewodniczanski Institute of Nuclear Physics, Krakow

M.-C. Chang, N. Kikuchi, Y. Mikami, T. Nagamine,

P. Schönmeier, A. Yamaguchi, and H. Yamamoto

Tohoku University, Sendai

P. Chang, Y. Chao, K.-F. Chen, W.-S. Hou, Y. B. Hsiung, Y.-J. Lee, C. Y. Lin,
S.-W. Lin, Y.-T. Shen, Y. T. Tsai, K. Ueno, C. C. Wang, M.-Z. Wang, and C.-H. Wu

Department of Physics, National Taiwan University, Taipei

B. G. Cheon

Chonnam National University, Kwangju

J. H. Choi, H. Ha, J. S. Kang, and E. Won

Korea University, Seoul

S.-K. Choi

Gyeongsang National University, Chinju

Y. Choi, Y. K. Choi, H. O. Kim, J. H. Kim, C. W. Park, and K. S. Park
Sungkyunkwan University, Suwon

A. Chuvikov, A. Garmash, D. Marlow, and T. Ziegler
Princeton University, Princeton, New Jersey 08544

M. Dash, D. Mohapatra, L. E. Piilonen, and Y. Yusa
Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061

M. Fujikawa, H. Hayashii, A. Imoto, S. U. Kataoka, K. Miyabayashi, and S. Noguchi
Nara Women's University, Nara

B. Golob and P. Križan
*University of Ljubljana, Ljubljana and
J. Stefan Institute, Ljubljana*

M. Grosse Perdekamp and R. Seidl
*University of Illinois at Urbana-Champaign, Urbana, Illinois 61801 and
RIKEN BNL Research Center, Upton, New York 11973*

T. Hara, D. Heffernan, and H. Miyake
Osaka University, Osaka

Y. Hasegawa, N. Satoyama, and N. Takada
Shinshu University, Nagano

K. Hoshina and O. Nitoh
Tokyo University of Agriculture and Technology, Tokyo

H. Ishino, H. R. Khan, A. Kibayashi, T. Mori, S. Ono, and Y. Watanabe
Tokyo Institute of Technology, Tokyo

M. Iwabuchi, Y. J. Kim, Y. Liu, T. R. Sarangi, and Y. Uchida
The Graduate University for Advanced Studies, Hayama

J. H. Kang, T. H. Kim, and Y.-J. Kwon

Yonsei University, Seoul

H. Kawai and E. Kurihara

Chiba University, Chiba

H. J. Kim and H. Park

Kyungpook National University, Taegu

S. K. Kim, J. Lee, S. E. Lee, and Heyoung Yang

Seoul National University, Seoul

R. Kumar, J. B. Singh, and N. Soni

Panjab University, Chandigarh

J. S. Lange

University of Frankfurt, Frankfurt

G. Leder, J. MacNaughton, F. Mandl, W. Mitaroff,

M. Pernicka, C. Schwanda, and L. Widhalm

Institute of High Energy Physics, Vienna

T. Matsumoto, T. Nakagawa, T. Seki, T. Sumiyoshi, and S. Yamamoto

Tokyo Metropolitan University, Tokyo

J. Mueller

University of Pittsburgh, Pittsburgh, Pennsylvania 15260

A. Murakami, A. Sugiyama, and S. Suzuki

Saga University, Saga

Y. Nagasaka

Hiroshima Institute of Technology, Hiroshima

E. Nakano, H. Sakaue, and Y. Teramoto

Osaka City University, Osaka

A. Ogawa

RIKEN BNL Research Center, Upton, New York 11973

S. Ogawa and H. Shibuya

Toho University, Funabashi

S. Okuno

Kanagawa University, Yokohama

H. Sakamoto

Kyoto University, Kyoto

J. Schümann and C. H. Wang

National United University, Miao Li

S. Stanič

University of Nova Gorica, Nova Gorica

Q. L. Xie, Y. Yuan, S. L. Zang, and C. C. Zhang

Institute of High Energy Physics, Chinese Academy of Sciences, Beijing

Y. Yamashita

Nippon Dental University, Niigata

L. M. Zhang and Z. P. Zhang

University of Science and Technology of China, Hefei

Abstract

The $D_s^\pm \rightarrow K^\pm K^\mp \pi^\pm$ absolute branching fraction is measured using $e^+e^- \rightarrow D_s^{*\pm} D_{s1}^\mp$ (2536) events collected by the Belle detector at the KEKB e^+e^- asymmetric energy collider. Using the ratio of yields when either the D_{s1} or D_s^* is fully reconstructed, we find $\mathcal{B}(D_s^\pm \rightarrow K^\pm K^\mp \pi^\pm) = (4.0 \pm 0.4(stat) \pm 0.4(sys))\%$.

PACS numbers: 14.40.Lb, 13.66.Bc, 13.25.Ft

Knowledge of D_s^+ meson¹ absolute branching fractions is important for normalization of many decays involving a D_s^+ in a final state. The poor accuracy of the branching fraction $\mathcal{B}(D_s^+ \rightarrow K^+ K^- \pi^+) = (5.2 \pm 0.9)\%$ [1] has been a systematic limitation for some precise measurements. In particular, the recent study of the CP violation in $B^0 \rightarrow D^{(*)\pm} \pi^\mp$ decays is restricted by the knowledge of the ratio of two amplitudes that determine the CP -asymmetry [2, 3]. The amplitude $B^0 \rightarrow D^{(*)+} \pi^-$ can be calculated from the branching fraction of $B^0 \rightarrow D_s^{(*)+} \pi^-$ decays assuming factorization. On the other hand, the factorization hypothesis can be tested by measuring the ratio of $B^0 \rightarrow D^{(*)-} \pi^+$ and $B^0 \rightarrow D^{(*)-} D_s^+$ decays. Both $\mathcal{B}(B^0 \rightarrow D_s^{(*)+} \pi^-)$ and $\mathcal{B}(B^0 \rightarrow D^{(*)-} D_s^+)$ measurements can be improved with better accuracy in D_s^+ absolute branching fractions.

Recently, the absolute branching fraction of $D_s^+ \rightarrow \phi(\rightarrow K^+ K^-) \pi^+$ was measured by the BaBar collaboration, which used partial and full reconstruction of $B \rightarrow D^{(*)} D_s^{(*)+}$ decays [4]. Another result obtained from a $\sqrt{s}$ -scan above $D_s^+ D_s^-$ threshold was presented by the CLEO-c collaboration [5].

In this paper we report on a measurement of the $D_s^+ \rightarrow K^+ K^- \pi^+$ branching fraction using two body $e^+ e^-$ -continuum annihilation into a $D_s^{*+} D_{s1}^- (2536)$ final state. The analysis is based on 552.3 fb^{-1} of data at the $\Upsilon(4S)$ resonance and nearby continuum, collected with the Belle detector [6] at the KEKB asymmetric energy storage ring [7].

I. METHOD

We use the partial reconstruction of the process $e^+ e^- \rightarrow D_s^{*+} D_{s1}^-$. In this analysis 4-momentum conservation allows us to infer the 4-momentum of the undetected part. The method used was described in Ref. [8] and applied to the measurement of the $e^+ e^- \rightarrow D^{(*)+} D^{(*)-}$ cross sections.

Here we reconstruct the process $e^+ e^- \rightarrow D_s^{*+} D_{s1}^-$ using two different tagging procedures. The first one (denoted as the D_{s1}^- tag) includes the full reconstruction of the D_{s1}^- meson via $D_{s1}^- \rightarrow \bar{D}^* K$ decay and observation of the photon from $D_s^{*+} \rightarrow D_s^+ \gamma$, while the D_s^+ is not reconstructed. The measured signal yield with the D_{s1}^- tag is proportional to the branching fractions of the reconstructed $\bar{D}^*$ modes. In the second procedure (denoted as the D_s^{*+} tag)

¹ Charge conjugation is implied through the paper.

we require full reconstruction of D_s^{*+} through $D_s^{*+} \rightarrow D_s^+ \gamma$ and observation of the kaon from $D_{s1}^- \rightarrow \bar{D}^* K$, but the $\bar{D}^*$ is not reconstructed. Since the D_s^+ meson is reconstructed in the channel of interest, $D_s^+ \rightarrow K^+ K^- \pi^+$, the signal yield measured with the D_s^{*+} tag is proportional to this D_s^+ branching fraction. The (efficiency-corrected) ratio of the two measured signal yields is equal to the ratio of well-known $\bar{D}^*$ branching fractions and the branching fraction of the D_s^+ .

In order to calculate reconstruction efficiencies and optimize event selection criteria, Monte Carlo (MC) signal events are generated and simulated using a GEANT-based full simulator, including initial state radiation (ISR), and assuming no form-factors for D_s^{*+} and D_{s1}^- mesons.

To identify the signal we study the mass recoiling against the reconstructed particle (or combination of particles) denoted as X . This recoil mass is defined as:

$$M_{\text{recoil}}(X) \equiv \sqrt{(E_{CM} - E_X)^2 - P_X^2}, \quad (1)$$

where E_X and P_X are the center-of-mass (CM) energy and momentum of X , respectively; E_{CM} is the CM beam energy. We expect a peak in the M_{recoil} distribution at the nominal mass of the recoil particle.

The resolution in M_{recoil} is $\sim 50 \text{ MeV}/c^2$ according to the MC, which is not sufficient to separate different final states, *e.g.* $D_s^+ D_{s1}^-$, $D_s^{*+} D_{s1}^-$ and non-resonant $D_s^+ \bar{D}^* K$. To disentangle the contribution of these final states we use another kinematic variable, the recoil mass difference ΔM_{recoil} :

$$\Delta M_{\text{recoil}}(D_{s1}^- \gamma) \equiv M_{\text{recoil}}(D_{s1}^-) - M_{\text{recoil}}(D_{s1}^- \gamma), \quad (2)$$

$$\Delta M_{\text{recoil}}(D_s^{*+} K) \equiv M_{\text{recoil}}(D_s^{*+}) - M_{\text{recoil}}(D_s^{*+} K). \quad (3)$$

In the D_{s1}^- tag procedure the signal events make a narrow peak in the $\Delta M_{\text{recoil}}(D_{s1}^- \gamma)$ distribution at the nominal $D_s^{*+} - D_s^+$ mass difference with a resolution of $\sigma \sim 5 \text{ MeV}/c^2$, dominated by the γ energy resolution according to the MC. The $\Delta M_{\text{recoil}}(D_s^{*+} K)$ spectrum peaks at the $D_{s1}^- - \bar{D}^*$ mass difference with a resolution of $\sigma \sim 2 \text{ MeV}/c^2$.

II. SELECTION

All primary charged tracks are required to be consistent with originating from the interaction point. Charged kaon candidates are identified using information from dE/dx

measurements in the central drift chamber, aerogel Cherenkov counters and time-of-flight system. No identification requirements are applied for pion candidates. K_S^0 candidates are reconstructed from $\pi^+\pi^-$ pairs with an invariant mass within $15 \text{ MeV}/c^2$ of the nominal K_S^0 mass. The distance between the two pion tracks at the K_S^0 vertex is required to be smaller than 1 cm, the flight distance in the plane perpendicular to the beam from the interaction point is required to be greater than 0.1 cm and the angle between the K_S^0 momentum direction and decay path in this plane should be smaller than 0.01 rad. Photons are reconstructed in the electromagnetic calorimeter as showers with energies above 50 MeV that are not associated with charged tracks. π^0 candidates are reconstructed by combining pairs of photons with invariant masses within $15 \text{ MeV}/c^2$ of the nominal π^0 mass.

D^0 candidates are reconstructed using five decay modes: $K^-\pi^+$, K^-K^+ , $K^-\pi^-\pi^+\pi^+$, $K_S^0\pi^+\pi^-$ and $K^-\pi^+\pi^0$. D^+ candidates are reconstructed using the $K^-\pi^+\pi^+$ channel. D_s^+ candidates are reconstructed using the $K^+K^-\pi^+$ decay mode. A $\pm 15 \text{ MeV}/c^2$ mass window is used for all D modes (approximately 2.5σ in each case) except for the $D^0 \rightarrow K^-\pi^-\pi^+\pi^+$ where $\pm 10 \text{ MeV}/c^2$ is applied and $D^0 \rightarrow K^-\pi^+\pi^0$ with $\pm 20 \text{ MeV}/c^2$ mass window. All D_s^+ , D^+ and D^0 candidates are subjected to a mass and vertex constrained fit to improve their momenta and thus the recoil mass resolution.

D^* candidates are selected via $D^{*+} \rightarrow D^0\pi^+$ and $D^{*0} \rightarrow D^0\pi^0$ decay modes with $\pm 2 \text{ MeV}/c^2$ mass window. D_s^{*+} candidates are reconstructed via $D_s^+\gamma$ channel with $\pm 10 \text{ MeV}/c^2$ mass window. D_{s1}^- is reconstructed in $\overline{D}^{*0}K^-$ and $D^{*-}K_S^0$ decay modes.

In the case of multiple candidates, the candidate with the minimum value of χ_{tot}^2 ($\chi_{tot}^2 = \chi_{M(D)}^2 + \chi_{M(D^*)}^2$ for D_{s1}^- tag and $\chi_{tot}^2 = \chi_{M(D_s^+)}^2$ for D_s^{*+} tag, respectively) is chosen. Each χ^2 is defined as the square of the ratio of the deviation of the measured mass from the nominal value and the corresponding resolution.

III. RECONSTRUCTION

A. D_{s1}^- tag

As the ratio of $D_{s1}^- \rightarrow \overline{D}^{*0}K^-$ and $D_{s1}^- \rightarrow D^{*-}K_S^0$ branching fractions is unknown, we perform the analysis for these two channels separately and average the results.

The $\overline{D}^{*0}K^-$ and $D^{*-}K_S^0$ mass spectra with a preselection requirement on $M_{\text{recoil}}(\overline{D}^*K) <$

3 GeV/c^2 are shown in Fig. 1. Clean peaks at the nominal D_{s1}^- mass without substantial combinatorial background are evident in both distributions. We fit these distributions with a sum of two Gaussians with a common mean and the background parametrized with a threshold function:

$$Bg(m) = (A \cdot m + B) \sqrt{m - M_{D^*}^{PDG} - M_K^{PDG}}. \quad (4)$$

The fit yields a Gaussian central value of $M = 2535.1 \pm 0.4 \text{ MeV}/c^2$ for $\bar{D}^{*0} K^-$ and $M = 2535.0 \pm 0.3 \text{ MeV}/c^2$ for $D^{*-} K_S^0$ channels, respectively, which is in good agreement with both the PDG and MC values.




FIG. 1: The distribution of invariant mass a) $\bar{D}^{*0} K^-$ and b) $D^{*-} K_S^0$. The solid curves represent the fit described in the text.

The signal and sideband regions for D_{s1}^- are defined as $|M(D_{s1}^-) - 2.5354| < 0.005 \text{ GeV}/c^2$ and $0.010 \text{ GeV}/c^2 < |M(D_{s1}^-) - 2.5354| < 0.015 \text{ GeV}/c^2$, respectively. The $M_{\text{recoil}}(D_{s1}^-)$ spectra are shown in Fig. 2. The significant excess in the region from 1.9 to 2.2 GeV/c^2 includes

$e^+e^- \rightarrow D_s^+ D_{s1}^-$ and $e^+e^- \rightarrow D_s^{*+} D_{s1}^-$ processes. The small excess around $2.3 \text{ GeV}/c^2$ could be interpreted as a contribution from the process $e^+e^- \rightarrow D_{sJ}^+ D_{s1}^-$. The excess around $2.5 \text{ GeV}/c^2$ corresponds to the excited $D_s^{**+} = D_{s1}^+, D_{s2}^+$ states.




FIG. 2: a) $M_{\text{recoil}}(D_{s1}^- \rightarrow \overline{D}^{*0} K^-)$ distribution; b) $M_{\text{recoil}}(D_{s1}^- \rightarrow D^{*-} K_S^0)$ distribution. Points with error bars correspond to the events in the signal region. The D_{s1}^- sideband contribution is shown as hatched histogram. The signal windows are indicated by the vertical lines.

We define the signal $M_{\text{recoil}}(D_{s1}^-)$ window as $|M_{\text{recoil}}(D_{s1}^-) - 2.1121| < 0.100 \text{ GeV}/c^2$ (Fig. 2). As the recoil mass resolution of $\sim 50 \text{ MeV}/c^2$ is not sufficient to resolve $e^+e^- \rightarrow D_s^+ D_{s1}^-$ and $e^+e^- \rightarrow D_s^{*+} D_{s1}^-$ processes, to disentangle them we reconstruct the γ from $D_s^{*+} \rightarrow D_s^+ \gamma$ decays. The reconstructed D_{s1}^- candidates from the signal $M_{\text{recoil}}(D_{s1}^-)$ region are combined with each photon in the event to calculate the recoil mass difference $\Delta M_{\text{recoil}}(D_{s1}^- \gamma)$.

To obtain the signal yield and disentangle the signal contribution from the non-resonant $D_s^{*+} \overline{D}^* K$ process, we perform fits to $\overline{D}^* K$ mass distributions in bins of $\Delta M_{\text{recoil}}(D_{s1}^- \gamma)$. The

$\Delta M_{\text{recoil}}(D_{s1}^- \gamma)$ bin width is chosen to be $4 \text{ MeV}/c^2$. The signal function is the sum of two Gaussians with the mass and width fixed to the corresponding MC values. The background is described with a threshold function (Eq. 4). The signal yield for each bin of the recoil mass difference for the $D_{s1}^- \rightarrow \bar{D}^{*0} K^-$ and $D_{s1}^- \rightarrow D^{*-} K_S^0$ channels are plotted in Fig. 3.




FIG. 3: The signal yields in bins of $\Delta M_{\text{recoil}}(D_{s1}^- \gamma)$: a) for the $D_{s1}^- \rightarrow \bar{D}^{*0} K^-$ channel; b) for the $D_{s1}^- \rightarrow D^{*-} K_S^0$ channel. The solid curves represent the fits described in the text.

The method used to extract the signal yield guarantees that the signal peak includes only those events that contain both a D_{s1}^- meson and a D_s^{*+} decaying to $D_s \gamma$. Thus, the only possible peaking background sources are final states with extra pion(s) (e.g. $e^+ e^- \rightarrow D_s^{*+} D_{s1}^- \pi^0$). The process with one extra π^0 is suppressed due to isospin conservation. The possible contribution of $e^+ e^- \rightarrow D_{s1}^- D_{sJ}^+(2460)$ followed by $D_{sJ}^+(2460) \rightarrow D_s^{*+} \pi^0$ decay, where the isospin violation occurs in the D_{sJ} decay, is suppressed by the recoil mass cut at the 4σ level. The processes with two pions are also suppressed at the $\sim 3.5\sigma$ level. Therefore, we conclude that the remaining background is negligibly small; the extracted yields are

dominated by the signal.

The distributions in Fig. 3 are fitted to the sum of a signal Gaussian (with the mass and width fixed to the MC values) and a linear background function. The fits yield 184 ± 19 signal events for $D_{s1}^- \rightarrow \bar{D}^{*0} K^-$ and 105 ± 14 events for $D_{s1}^- \rightarrow D^{*-} K_S^0$. To determine the reconstruction efficiency the same procedure is applied to the MC signal sample.

B. D_s^{*+} tag

In the D_s^{*+} tag, the D_s^{*+} of the process $e^+e^- \rightarrow D_s^{*+} D_{s1}^-$ is fully reconstructed. The $D_s^+ \gamma$ mass distribution is shown in Fig. 4 and fitted with the sum of a signal Gaussian and a linear background function. The obtained central value of the Gaussian $M = 2113.5 \pm 0.4 \text{ MeV}/c^2$ is somewhat larger than the PDG value, but in good agreement with the MC expectation, which shows a similar shift. The signal region is chosen to be $|M(D_s^+ \gamma) - 2.1121| < 0.010 \text{ GeV}/c^2$ while the sidebands are defined as $0.015 \text{ GeV}/c^2 < |M(D_s^+ \gamma) - 2.1121| < 0.025 \text{ GeV}/c^2$.




FIG. 4: The $D_s^+ \gamma$ invariant mass distribution. The solid curve represents the fit described in the text.

The $M_{\text{recoil}}(D_s^{*+})$ spectrum for the signal region is shown in Fig. 5. A wide enhancement around $2.1 \text{ GeV}/c^2$ is formed by overlapping D_s^- and D_s^{*-} signals. The signal process $e^+e^- \rightarrow D_s^{*+} D_{s1}^-$ contributes to the bump around $2.55 \text{ GeV}/c^2$ together with the process

$e^+e^- \rightarrow D_s^{*+}D_{s2}^-$. We define the signal region by the requirement $|M_{\text{recoil}}(D_s^{*+}) - 2.5354| < 0.100 \text{ GeV}/c^2$ ($\sim 2\sigma$).




FIG. 5: The $M_{\text{recoil}}(D_s^{*+})$ distribution. The events in the D_s^{*+} signal region are shown as points with error bars. The D_s^{*+} sidebands are superimposed as hatched histogram. The $M_{\text{recoil}}(D_s^{*+})$ signal window is indicated by vertical lines.

To extract the $e^+e^- \rightarrow D_s^{*+}D_{s1}^-$ signal yield for D_s^{*+} tag, we perform a fit to the $D_s^+\gamma$ mass distributions in bins of $\Delta M_{\text{recoil}}(D_s^{*+}K)$. The signal function is a Gaussian with its parameters fixed to the corresponding MC values. The background is parametrized by a linear function. The yields returned by the fits for $D_{s1}^- \rightarrow \bar{D}^{*0}K^-$ and $D_{s1}^- \rightarrow D^{*-}K_S^0$ channels are plotted in Fig. 6. We fit the resulting distribution with the sum of a signal Gaussian (with fixed mass and width) and a threshold function (Eq. 4). The fits yield 267 ± 18 signal events for the $\bar{D}^{*0}K^-$ channel and 45 ± 8 events for the $D^{*-}K_S^0$ channel.

IV. SYSTEMATIC ERRORS

The systematic uncertainties are summarized in Table I.

Different production and helicity angular distributions result in different ratios of reconstruction efficiencies for the two tagging procedures. A potential source of significant systematic error are the uncertainties, associated with reconstruction efficiencies of the kaon from the D_{s1}^- decay and the photon from the D_s^{*+} , which strongly depend on the mother




FIG. 6: The $e^+e^- \rightarrow D_s^{*+}D_{s1}^-$ yields for D_s^{*+} tag in bins of $\Delta M_{\text{recoil}}(D_s^{*+}K)$: a) for the $D_{s1}^- \rightarrow \bar{D}^{*0}K^-$ channel; b) for the $D_{s1}^- \rightarrow D^{*-}K_S^0$ channel. The solid curves show the fits described in the text.

particle polarizations. As the kaon and photon are, however, reconstructed in both tags, the systematic uncertainties related to the reconstruction of these particles cancel out. The remaining systematic uncertainty in the efficiency ratio from the production angle θ_{prod} (polar angle of the momentum of D_{s1}^- meson) is estimated from MC simulation by comparing the regions $|\cos(\theta_{\text{prod}})| > 0.5$ and $|\cos(\theta_{\text{prod}})| < 0.5$ and found to be small (0.4%).

The systematic error arising from unknown $\bar{D}^*$ polarizations contributes only to the uncertainty in the efficiency of D_{s1}^- tag and does not cancel out in the efficiency ratio. This contribution is estimated from MC assuming different $\bar{D}^*$ polarizations.

The efficiency of the requirement on recoil mass depends on the fraction of events with an ISR photon as well as on the recoil mass window cut. The error from the ISR photon is conservatively estimated by changing the fraction of events containing ISR photons in

TABLE I: Contributions to the systematic error.

Source	Estimated error, [%]
Production angle	0.4
$\overline{D}^*$ polarization	3.8
ISR	2.3
Recoil mass cut	4.8
Fit	4.3
Tracking/ π^0 reconstr. efficiency	3.0
$\mathcal{B}(\overline{D}^{(*)})$	5.2
MC statistics	2.3
Total	10.1

the whole MC sample by 30%. The resulting relative shift in $\mathcal{B}(D_s^+)$ is found to be 2.3%. The error from the recoil mass cut is estimated by varying the recoil mass window width in the range of $80 - 120 \text{ MeV}/c^2$. The difference in the calculated value of $\mathcal{B}(D_s^+)$ (4.8%) is conservatively considered as the systematic error.

The fit systematics is estimated assuming different shapes for the signal (single and double Gaussians) and for the background (different order polynomials). The difference in the resulting branching fraction is found to be 4.3% and is treated as the systematic uncertainty in the fit.

Other contributions come from the uncertainties in the track/ π^0 reconstruction efficiencies, $\overline{D}^{(*)}$ branching fractions and MC statistics.

All of the systematic contributions described above are added in quadrature to give 10.1%.

V. RESULTS

Using the measured signal yields $N(D_s^{*+})$ and $N(D_{s1}^-)$ with D_s^{*+} and D_{s1}^- tags, respectively, and taking into account the efficiency ratio ($\frac{\epsilon(D_{s1}^-)}{\epsilon(D_s^{*+})} = 0.50$ for $\overline{D}^{*0} K^-$ and $\frac{\epsilon(D_{s1}^-)}{\epsilon(D_s^{*+})} = 1.65$ for $D^{*-} K_S^0$ channels, respectively), we obtain the D_s^+ absolute branching fraction:

$$\mathcal{B}(D_s^+ \rightarrow K^+ K^- \pi^+) = \frac{N(D_s^{*+})}{N(D_{s1}^-)} \cdot \frac{\epsilon(D_{s1}^-)}{\epsilon(D_s^{*+})} \cdot \mathcal{B}(\overline{D}^{(*)}), \quad (5)$$

where $\mathcal{B}(\overline{D}^{(*)})$ is the product of $\overline{D}^*$ branching fraction and those of sub-decays. Separate calculations for the $\overline{D}^{*0}K^-$ and $D^{*-}K_S^0$ channels yield $\mathcal{B}(D_s^+ \rightarrow K^+K^-\pi^+)$ of $(4.01 \pm 0.47(stat))\%$ and $(3.84 \pm 0.83(stat))\%$, respectively.

VI. SUMMARY

In summary, we report a measurement of the $D_s^+ \rightarrow K^+K^-\pi^+$ branching fraction using a new method of double tag partial reconstruction. The branching fraction is measured separately in two channels $e^+e^- \rightarrow D_s^{*+}D_{s1}^-(\rightarrow \overline{D}^{*0}K^-)$ and $e^+e^- \rightarrow D_s^{*+}D_{s1}^-(\rightarrow D^{*-}K_S^0)$. The average value is $\mathcal{B}(D_s^+ \rightarrow K^+K^-\pi^+) = (4.0 \pm 0.4(stat) \pm 0.4(sys))\%$.

-
- [1] W.-M. Yao *et al.* (Particle Data Group), J. Phys. G **33**, 1 (2006).
 - [2] F. J. Ronga *et al.* (Belle collaboration), Phys. Rev. D **73**, 092003 (2006).
 - [3] B. Aubert *et al.* (BaBar collaboration), Phys. Rev. D **71**, 112003 (2005).
 - [4] B. Aubert *et al.* (BaBar collaboration), Phys. Rev. D **74**, 031103 (2006).
 - [5] N. E. Adam *et al.* (Cleo-c collaboration), hep-ex/0607079.
 - [6] A. Abashian *et al.* (Belle collaboration), Nucl. Instr. and Meth. A **479**, 117 (2002).
 - [7] S. Kurokawa and E. Kikutani, Nucl. Instr. and Meth. A **499**, 1 (2003), and other papers included in this volume.
 - [8] T. Uglov, *et al.* (Belle collaboration), Phys. Rev. D **70**, 071101 (2004).